

Suzuki, M., et al. 2013. Deer herbivory as an important driver of divergence of ground vegetation communities in temperate forests. – *Oikos* 122: 104–110.

Appendix 1

Table A1. The most abundant (major) ground-layer vegetation species in mixed temperate forests of Boso peninsula, Japan.

Species	Abbreviation	Family	Life form
<i>Achyranthes bidentata</i>	Acbi	Amaranthaceae	Forb
<i>Alpinia japonica</i>	Alja	Zingiberaceae	Forb
<i>Arachniodes aristata</i>	Arar	Dryopteridaceae	Fern
<i>Arachniodes sporadosora</i>	Arsp	Dryopteridaceae	Fern
<i>Arachniodes standishii</i>	Arst	Dryopteridaceae	Fern
<i>Ardisia japonica</i>	Arja	Myrsinaceae	Shrub
<i>Aucuba japonica</i>	Auja	Cornaceae	Shrub
<i>Boenninghausenia albiflora</i>	Boal	Rutaceae	Fern
<i>Carex lenta</i>	Cale	Cyperaceae	Grass
<i>Carex morrowii</i>	Camo	Cyperaceae	Grass
<i>Carex multifolia</i>	Camu	Cyperaceae	Grass
<i>Castanopsis sieboldii</i>	Cssi	Fagaceae	Tree
<i>Cinnamomum tenuifolium</i>	Ciin	Lauraceae	Tree
<i>Damnacanthus indicus</i>	Dain	Rubiaceae	Shrub
<i>Dryopteris erythrosora</i>	Drer	Dryopteridaceae	Fern
<i>Eurya japonica</i>	Euja	Theaceae	Tree
<i>Iris japonica</i>	Irja	Iridaceae	Forb
<i>Lindera umbellata</i>	Lium	Lauraceae	Tree
<i>Maesa japonica</i>	Maja	Myrsinaceae	Shrub
<i>Neolitsea sericea</i>	Nese	Lauraceae	Tree
<i>Ophiopogon japonicus</i>	Oppj	Liliaceae	Forb
<i>Oplismenus undulatifolius</i>	Opun	Poaceae	Grass
<i>Osmunda japonica</i>	Osja	Osmundaceae	Fern
<i>Pertya scandens</i>	Pesc	Asteraceae	Shrub
<i>Piper kadsura</i>	Pika	Piperaceae	Forb
<i>Quercus acuta</i>	Quac	Fagaceae	Tree
<i>Rubus buergeri</i>	Rubu	Rosaceae	Vine
<i>Selaginella remotifolia</i>	Sere	Selaginellaceae	Fern
<i>Trachelospermum asiaticum</i>	Tras	Apocynaceae	Vine