

Yoshikawa, T., Kawakami, K. and Masaki, T. 2019. Allometric scaling of seed retention time in seed dispersers and its application to estimation of seed dispersal potentials of theropod dinosaurs. – Oikos doi: 10.1111/oik.05827

Appendix 1

List of the publications from which we extracted mean seed retention times (SRTs) of animals for our analysis.

Agami, M. and Waisel, Y. 1988. The role of fish in distribution and germination of seeds of the submerged macrophytes *Najas marina* L. and *Ruppia maritima* L. – Oecologia 76: 83–88.

Anderson, J. T. et al. 2011. Extremely long-distance seed dispersal by an overfished Amazonian frugivore. – Proc. R. Soc. B 278: 3329–3335.

Blake, S. et al. 2012. Seed dispersal by Galapagos tortoises. – J. Biogeogr. 39: 1961–1972.

Brochet, A.-L. et al. 2010. Endozoochory of Mediterranean aquatic plant seeds by teal after a period of desiccation: determinants of seed survival and influence of retention time on germinability and viability. – Aquat. Bot. 93: 99–106.

Campos-Arceiz, A. et al. 2008. Behavior rather than diet mediates seasonal differences in seed dispersal by Asian elephants. – Ecology 89: 2684–2691.

Campos-Arceiz, A. et al. 2012. Asian tapirs are no elephants when it comes to seed dispersal. – Biotropica 44: 220–227.

Castilla, A. M. 2000. Does passage time through the lizard *Podarcis lilfordi*'s guts affect germination performance in the plant *Withania frutescens*? – Acta Oecol. 21: 119–124.

Chang, S.-Y. et al. 2012. Frugivory by Taiwan barbets (*Megalaima nuchalis*) and the effects of

- deinhibition and scarification on seed germination. – *Can. J. Zool.* 90: 640–650.
- Chang, N.-C. et al. 2016. Effects of dietary fiber on gut retention time in captive *Macaca cyclopis*, *Macaca fascicularis*, *Hylobates lar* and *Pongo pygmaeus* and the germination of ingested seeds. – *Int. J. Primatol.* 37: 671–687.
- Charalambidou, I. et al. 2003. Effect of ingestion by five avian dispersers on the retention time, retrieval and germination of *Ruppia maritima* seeds. – *Funct. Ecol.* 17: 747–753.
- Charalambidou, I. et al. 2005. Digestive plasticity in Mallard ducks modulates dispersal probabilities of aquatic plants and crustaceans. – *Funct. Ecol.* 19: 513–519.
- Clout, M. N. and Tilley, J. A. V. 1992. Germination of miro (*Prumnopitys ferruginea*) seeds after consumption by New Zealand pigeons (*Hemiphaga novaeseelandiae*). – *N. Z. J. Bot.* 30: 25–28.
- Cosyns, E. et al. 2005. Germination success of temperate grassland species after passage through ungulate and rabbit guts. – *J. Ecol.* 93: 353–361.
- D'hondt, B. and Hoffmann, M. 2011. A reassessment of the role of simple seed traits in mortality following herbivore ingestion. – *Plant Biol.* 13: 118–124.
- Dinerstein, E. and Wemmer, C. M. 1988. Fruits Rhinoceros eat: dispersal of *Trewia nudiflora* (Euphorbiaceae) in lowland Nepal. – *Ecology* 69: 1768–1774.
- Elfström, M. et al. 2013. Gut retention time in captive brown bears *Ursus arctos*. – *Wildl. Biol.* 19: 317–324.
- Figuerola, J. et al. 2010. Internal dispersal of seeds by waterfowl: effect of seed size on gut passage time and germination patterns. – *Naturwissenschaften* 97: 555–565.
- Fukui, A. 2003. Relationship between seed retention time in bird's gut and fruit characteristics. – *Ornithol. Sci.* 2: 41–48.
- Garcia-Alvarez, A. et al. 2015. Internal transport of alien and native plants by geese and ducks: an experimental study. – *Freshw. Biol.* 60: 1316–1329.
- Gardener, C. J. et al. 1993. Passage of legume and grass seeds through the digestive tract of cattle and their survival in faeces. – *J. Appl. Ecol.* 30: 63–74.

- Gervais, J. A. et al. 1998. The potential for seed dispersal by the banana slug (*Ariolimax columbianus*). – Am. Midl. Nat. 140: 103–110.
- Giordani, L. et al. 2015. The domestic goat as a potential seed disperser of *Mimosa luisana* (Leguminosae, Mimosoideae) in the Tehuacan-Cuicatlan Valley, Mexico. – J. Trop. Ecol. 31: 91–94.
- Graae, B. J. et al. 2004. An experimental evaluation of the arctic fox (*Alopex lagopus*) as a seed disperser. – Arct. Antarct. Alp. Res. 36: 468–473.
- Graham, C. H. et al. 1995. Seed dispersal effectiveness by two bulbuls on *Maesa lanceolata*, an African montane forest tree. – Biotropica 27: 479–486.
- Green, A. K. et al. 2009. Directed dispersal of mistletoe (*Plicosepalus acaciae*) by Yellow-vented Bulbuls (*Pycnonotus xanthopygos*). – J. Ornithol. 150: 167–173.
- Guix, J. C. and Ruiz, X. 1997. Weevil larvae dispersal by guans in southeastern Brazil. – Biotropica 29: 522–525.
- Guzman, A. and Stevenson, P. R. 2011. A new hypothesis for the importance of seed dispersal in time. – Rev. Biol. Trop. 59: 1795–1803.
- Harich, F. K. et al. 2016. Seed dispersal potential of Asian elephants. – Acta Oecol. 77: 144–151.
- Hickey, J. R. et al. 1999. An evaluation of a mammalian predator, *Martes americana*, as a disperser of seeds. – Oikos 87: 499–508.
- Holbrook, K. M. and Smith, T. B. 2000. Seed dispersal and movement patterns in two species of *Ceratogymna* hornbills in a West African tropical lowland forest. – Oecologia 125: 249–257.
- Holbrook, K. M. et al. 2007. Using toucan-generated dispersal models to estimate seed dispersal in Amazonian Ecuador. – In: Dennis, A. J. et al. (eds), Seed dispersal: theory and its application in a changing world. pp. 300–321.
- Horn, M. H. 1997. Evidence for dispersal of fig seeds by the fruit-eating characid fish *Brycon guatemalensis* Regan in a Costa Rican tropical rain forest. – Oecologia 109: 259–264.
- Janzen, D. H. 1981b. *Enterolobium cyclocarpum* seed passage rate and survival in horses, Costa Rican Pleistocene seed dispersal agents. – Ecology 62: 593–601.

- Janzen, D. H. 1981a. Digestive seed predation by a Costa Rican Baird's tapir. – *Biotropica* 13: 59–63.
- Janzen, D. H. 1982. Differential seed survival and passage rates in cows and horses, surrogate Pleistocene dispersal agents. – *Oikos* 38: 150–156.
- Jerozolinski, A. et al. 2009. Are tortoises important seed dispersers in Amazonian forests? – *Oecologia* 161: 517–528.
- Khamcha, D. et al. 2014. Behavioral and social structure effects on seed dispersal curves of a forest-interior bulbul (*Pycnonotidae*) in a tropical evergreen forest. – *Biotropica* 46: 294–301.
- Kleyheeg, E. et al. 2015. Bird-mediated seed dispersal: reduced digestive efficiency in active birds modulates the dispersal capacity of plant seeds. – *Oikos* 124: 899–907.
- Lambert, J. E. 2002. Digestive retention times in forest guenons (*Cercopithecus* spp.) with reference to chimpanzees (*Pan troglodytes*). – *Int. J. Primatol.* 23: 1169–1185.
- Lara, C. et al. 2009. Provenance, guts and fate: field and experimental evidence in a host-mistletoe-bird system. – *Ecoscience* 16: 399–407.
- Lehouck, V. et al. 2009. Complementary seed dispersal by three avian frugivores in a fragmented Afromontane forest. – *J. Veg. Sci.* 20: 1110–1120.
- Lenz, J. et al. 2011. Seed-dispersal distributions by trumpeter hornbills in fragmented landscapes. – *Proc. R. Soc. B* 278: 2257–2264.
- Levey, D. J. and Grajal, A. 1991. Evolutionary implications of fruit-processing limitations in cedar waxwings. – *Am. Nat.* 138: 171–189.
- Link, A. and Di Fiore, A. 2006. Seed dispersal by spider monkeys and its importance in the maintenance of neotropical rain-forest diversity. – *J. Trop. Ecol.* 22: 235–246.
- Mancilla-Leyton, J. M. et al. 2011. Plant-ungulate interaction: goat gut passage effect on survival and germination of Mediterranean shrub seeds. – *J. Veg. Sci.* 22: 1031–1037.
- Mancilla-Leyton, J. M. et al. 2013. Effects of rabbit gut passage on seed retrieval and germination of three shrub species. – *Basic Appl. Ecol.* 14: 585–592.
- Manzano, P. et al. 2005. Sheep gut passage and survival of Mediterranean shrub seeds. – *Seed Sci.*

Res. 15: 21–28.

- Martinez, B. T. and Razafindratsima, O. H. 2014. Frugivory and seed dispersal patterns of the red-ruffed lemur, *Varecia rubra*, at a forest restoration site in Masoala National Park, Madagascar. – *Folia Primatol.* 85: 228–243.
- Meehan, H. J. et al. 2003. Retention time of beads in captive Torres imperial pigeons, *Ducula spilorrhoa spilorrhoa*. – *Notornis* 50: 167–168.
- Michael, P. J. et al. 2006. Sheep rumen digestion and transmission of weedy *Malva parviflora* seeds. – *Aust. J. Exp. Agric.* 46: 1251–1256.
- Mise, Y. et al. 2016. Comparing methods of acquiring mammalian endozoochorous seed dispersal distance distributions. – *Ecol. Res.* 31: 881–889.
- Mokotjomela, T. M. et al. 2015. The potential for birds to disperse the seeds of *Acacia cyclops*, an invasive alien plant in South Africa. – *Ibis* 157: 449–458.
- Morales, J. M. et al. 2013. Frugivore behavioural details matter for seed dispersal: a multi-species model for Cantabrian thrushes and trees. – *PLoS One* 8: e65216.
- Moses, K. L. and Semple, S. 2011. Primary seed dispersal by the black-and-white ruffed lemur (*Varecia variegata*) in the Manombo forest, south-east Madagascar. – *J. Trop. Ecol.* 27: 529–538.
- Mouissie, A. M. et al. 2005. Ecological correlates of seed survival after ingestion by fallow deer. – *Funct. Ecol.* 19: 284–290.
- Murphy, S. R. et al. 1993. Differential passage time of mistletoe fruits through the gut of honeyeaters and flowerpeckers: effects on seedling establishment. – *Oecologia* 93: 171–176.
- Murray, K. 1988. Avian seed dispersal of three Neotropical gap-dependent plants. – *Ecol. Monogr.* 58: 271–298.
- Murray, K. et al. 1993. The Influence of seed packaging and fruit color on feeding preferences. – *Vegetatio* 108: 217–226.
- Nakashima, Y. and Sukor, J. A. 2010. Importance of common palm civets (*Paradoxurus hermaphroditus*) as a long-distance disperser for large-seeded plants in degraded forests. –

Tropics 18: 221–229.

- Nogales, M. et al. 1998. Shrikes, lizards and *Lycium intricatum* (Solanaceae) fruits: a case of indirect seed dispersal on an oceanic island (Alegranza, Canary Islands). – J. Ecol. 86: 866–871.
- Nogales, M. et al. 2001. Ecological and biogeographical implications of yellow-legged gulls (*Larus cachinnans* Pallas) as seed dispersers of *Rubia fruticosa* Ait. (Rubiaceae) in the Canary Islands. – J. Biogeogr. 28: 1137–1145.
- Otani, T. 2004. Effects of macaque ingestion on seed destruction and germination of a fleshy-fruited tree, *Eurya emarginata*. – Ecol. Res. 19: 495–501.
- Petre, C.-A. et al. 2015. Western lowland gorilla seed dispersal: are seeds adapted to long gut retention times? – Acta Oecol. 67: 59–65.
- Pollux, B. J. A. et al. 2006. The effect of seed morphology on the potential dispersal of aquatic macrophytes by the common carp (*Cyprinus carpio*). – Freshw. Biol. 51: 2063–2071.
- Pollux, B. J. A. et al. 2007. Consequences of intraspecific seed-size variation in *Sparganium emersum* for dispersal by fish. – Funct. Ecol. 21: 1084–1091.
- Power, M. L. and Myers, E. W. 2009. Digestion in the common marmoset (*Callithrix jacchus*), a gummivore–frugivore. – Am. J. Primatol. 71: 957–963.
- Ramirez, M. and Ornelas, J. F. 2009. Germination of *Psittacanthus schiedeanus* (mistletoe) seeds after passage through the gut of cedar waxwings and grey silky-flycatchers. – J. Torrey Bot. Soc. 136: 322–331.
- Razafindratsima, O. H. et al. 2014. Patterns of movement and seed dispersal by three lemur species. – Am. J. Primatol. 76: 84–96.
- Razanamandranto, S. et al. 2004. Effects of gut treatment on recovery and germinability of bovine and ovine ingested seeds of four woody species from the Sudanian savanna in West Africa. – Flora 199: 389–397.
- Rick, C. M. and Bowman, R. I. 1961. Galapagos tomatoes and tortoises. – Evolution 15: 407–417.
- Righini, N. et al. 2004. Effect of different primate species on germination of *Ficus* (*Urostigma*)

- seeds. – *Zoo Biol.* 23: 273–278.
- Santamaría, L. et al. 2007. Predicting spatial patterns of plant recruitment using animal-displacement kernels. – *PLoS One* 2: e1008.
- Sato, H. 2009. Gut passage time and size of swallowed seeds in the common brown lemur and the mongoose lemur. – *Primate Res.* 25: 45–54.
- Sekar, N. et al. 2015. In the elephant's seed shadow: the prospects of domestic bovids as replacement dispersers of three tropical Asian trees. – *Ecology* 96: 2093–2105.
- Setlalekgomo, M. R. and Sesinyi, K. 2014. Seed dispersal by serrated tortoises (*Psammobates oculiferus*) and the effect of their gut passage on seed germination. – *Sci. J. Anim. Sci.* 3: 252–257.
- Shi, T.-T. et al. 2015. Effects of frugivorous birds on seed retention time and germination in Xishuangbanna, southwest China. – *Zool. Res.* 36: 241–247.
- Soto-Gamboa, M. and Bozinovic, F. 2002. Fruit-disperser interaction in a mistletoe-bird system: a comparison of two mechanisms of fruits processing on seed germination. – *Plant Ecol.* 159: 171–174.
- Spiegel, O. and Nathan, R. 2007. Incorporating dispersal distance into the disperser effectiveness framework: frugivorous birds provide complementary dispersal to plants in a patchy environment. – *Ecol. Lett.* 10: 718–728.
- Stanley, M. C. and Lill, A. 2002a. Does seed packaging influence fruit consumption and seed passage in an avian frugivore? – *Condor* 104: 136–145.
- Stanley, M. C. and Lill, A. 2002b. Avian fruit consumption and seed dispersal in a temperate Australian woodland. – *Austral Ecol.* 27: 137–148.
- Stevenson, P. R. 2000. Seed dispersal by woolly monkeys (*Lagothrix lagothricha*) at Tinigua National Park, Colombia: dispersal distance, germination rates and dispersal quantity. – *Am. J. Primatol.* 50: 275–289.
- Sun, C. et al. 1997. Effectiveness of three turacos as seed dispersers in a tropical montane forest. – *Oecologia* 112: 94–103.

- Tewksbury, J. J. et al. 2008. Costs and benefits of capsaicin-mediated control of gut retention in dispersers of wild chilies. – *Ecology* 89: 107–117.
- Tsuji, Y. et al. 2010. Effects of the physical characteristics of seeds on gastrointestinal passage time in captive Japanese macaques. – *J. Zool.* 280: 171–176.
- Tsuji, Y. et al. 2011. Gastrointestinal passage time of seeds ingested by captive Japanese martens *Martes melampus*. – *Acta Theriol.* 56: 353–357.
- Tsuji, Y. et al. 2015. Effects of food intake on digesta passage time in captive Japanese martens (*Martes melampus*) and implications for endozoochorous seed dispersal. – *Mammal Study* 40: 13–18.
- Tuerke, M. et al. 2010. Seed consumption and dispersal of ant-dispersed plants by slugs. – *Oecologia* 163: 681–693.
- Valido, A. and Nogales, M. 2003. Digestive ecology of two omnivorous Canarian lizard species (*Gallotia*, Lacertidae). – *Amphib.-Reptil.* 24: 331–344.
- Varela, O. and Bucher, E. H. 2006. Passage time, viability, and germination of seeds ingested by foxes. – *J. Arid Environ.* 67: 566–578.
- Wahaj, S. A. et al. 1998. Control of gut retention time by secondary metabolites in ripe *Solanum* fruits. – *Ecology* 79: 2309–2319.
- Waibel, A. et al. 2013. Does a giant tortoise taxon substitute enhance seed germination of exotic fleshy-fruited plants? – *J. Plant Ecol.* 6: 57–63.
- Weerasinghe, U. R. et al. 1999. A record of the food retention time of the Asiatic elephant, *Elephas maximus*. – *Mammal Study* 24: 115–119.
- Westcott, D. A. and Graham, D. L. 2000. Patterns of movement and seed dispersal of a tropical frugivore. – *Oecologia* 122: 249–257.
- Westcott, D. A. et al. 2005. Incorporating patterns of disperser behaviour into models of seed dispersal and its effects on estimated dispersal curves. – *Oecologia* 146: 57–67.
- Westcott, D. A. et al. 2008. Cassowary dispersal of the invasive pond apple in a tropical rainforest: the contribution of subordinate dispersal modes in invasion. – *Divers. Distrib.* 14: 432–439.

- Williams, P. A. et al. 2000. Small mammals as potential seed dispersers in New Zealand. – *Austral Ecol.* 25: 523–532.
- Willson, M. F. 1989. Gut retention times of experimental pseudoseeds by emus. – *Biotropica* 21: 210–213.
- Wongsriphuek, C. et al. 2008. Dispersal of wetland plant seeds by mallards: influence of gut passage on recovery, retention and germination. – *Wetlands* 28: 290–299.
- Wotton, D. M. and Kelly, D. 2012. Do larger frugivores move seeds further? Body size, seed dispersal distance, and a case study of a large, sedentary pigeon. – *J. Biogeogr.* 39: 1973–1983.
- Yumoto, T. 1999. Seed dispersal by Salvin's curassow, *Mitu salvini* (Cracidae), in a tropical forest of Colombia: direct measurements of dispersal distance. – *Biotropica* 31: 654–660.
- Yumoto, T. et al. 1999. Estimation of the retention times and distances of seed dispersed by two monkey species, *Alouatta seniculus* and *Lagothrix lagotricha*, in a Colombian forest. – *Ecol. Res.* 14: 179–191.
- Zhou, Y.-B. et al. 2008. Frugivory and seed dispersal by a small carnivore, the Chinese ferret-badger, *Melogale moschata*, in a fragmented subtropical forest of central China. – *For. Ecol. Manag.* 255: 1595–1603.