

**OIK-06919**

Helsen, K., Van Cleemput, E., Bassi, L., Graae, B. J., Somers, B., Blonder, B. and Honnay, O. 2020. Inter- and intraspecific trait variation shape multidimensional trait overlap between two plant invaders and the invaded communities. – Oikos doi: 10.1111/oik.06919

## Appendix 1–11

Appendices 6 and 10 are supplied as separate .gif files

**Appendix 1. All species encountered in the sampled plots of the *Impatiens glandulifera* and *Rosa rugosa* datasets.** For each species indicated if it occurred in high abundance (HA) for at least one plot (in which case the traits were measured on field data) and whether it occurred in control plots (NAT<sub>C</sub>) and/or invaded plots (NAT<sub>I</sub>).

Species	<i>I. glandulifera</i> dataset			<i>R. rugosa</i> dataset		
	HA	NAT <sub>C</sub>	NAT <sub>I</sub>	HA	NAT <sub>C</sub>	NAT <sub>I</sub>
<i>Acer pseudoplatanus</i>			x		x	
<i>Achillea millefolium</i>		x			x	x
<i>Aegopodium podagraria</i>	x	x	x			
<i>Agrostis stolonifera</i>	x	x			x	x
<i>Alchemilla sp.</i>			x			
<i>Alliaria petiolata</i>			x			
<i>Allium vineale</i>					x	
<i>Alopecurus pratensis</i>	x	x	x			
<i>Ammophila arenaria</i>				x	x	x
<i>Angelica sylvestris</i>		x				
<i>Anthriscus sylvestris</i>	x	x				
<i>Arabis hirsuta</i>					x	x
<i>Arenaria serpyllifolia</i>					x	x
<i>Arrhenatherum elatius</i>		x	x	x	x	x
<i>Artemisia vulgaris</i>		x				
<i>Bellis perennis</i>					x	
<i>Betula pendula</i>		x				
<i>Brassica rapa</i>			x			
<i>Bromus hordeaceus</i>					x	x
<i>Bromus sterilis</i>	x	x	x	x	x	x
<i>Bryonia cretica</i>					x	
<i>Cakile maritima</i>					x	
<i>Calamagrostis canescens</i>	x	x	x			
<i>Cardamine flexuosa</i>			x			
<i>Carduus crispus</i>	x		x			
<i>Carex arenaria</i>				x	x	x
<i>Carex hirta</i>				x		x
<i>Carlina vulgaris</i>					x	x
<i>Cerastium arvense</i>					x	x
<i>Cerastium fontanum</i>					x	x
<i>Cerastium semidecandrum</i>					x	x
<i>Chaerophyllum temulum</i>			x		x	x
<i>Chamerion angustifolium</i>	x	x	x			
<i>Chenopodium album</i>	x		x			
<i>Circaeа lutetiana</i>			x			
<i>Cirsium arvense</i>	x	x	x		x	

<i>Cirsium helenioides</i>		x				
<i>Cirsium oleraceum</i>	x	x				
<i>Cirsium vulgare</i>				x	x	
<i>Claytonia perfoliata</i>				x	x	
<i>Clematis vitalba</i>				x		
<i>Convolvulus arvensis</i>				x		
<i>Convolvulus sepium</i>	x	x	x			
<i>Convolvulus soldanella</i>				x	x	
<i>Cotoneaster horizontalis</i>				x	x	x
<i>Crataegus monogyna</i>			x			x
<i>Crepis capillaris</i>				x	x	x
<i>Cynoglossum officinale</i>				x	x	x
<i>Cynosurus cristatus</i>			x			
<i>Dactylis glomerata</i>	x	x		x	x	x
<i>Deschampsia cespitosa</i>	x	x	x			
<i>Deschampsia flexuosa</i>		x	x			
<i>Diplotaxis tenuifolia</i>					x	x
<i>Dipsacus pilosus</i>	x		x			
<i>Dryopteris filix-mas</i>		x				
<i>Elymus athericus</i>					x	x
<i>Elytrigia atherica</i>				x		x
<i>Elytrigia juncea</i>				x	x	x
<i>Epilobium hirsutum</i>	x	x	x			
<i>Epilobium sp.</i>	x	x	x			
<i>Epipactis helleborine</i>		x			x	
<i>Equisetum arvense</i>	x		x			
<i>Equisetum palustre</i>		x				
<i>Erigeron acris</i>						x
<i>Erigeron canadensis</i>					x	x
<i>Erodium cicutarium</i>					x	
<i>Erophila verna</i>					x	x
<i>Euonymus europaeus</i>				x		x
<i>Fallopia convolvulus</i>			x			
<i>Festuca arenaria</i>				x	x	x
<i>Festuca arundinacea</i>		x				
<i>Festuca rubra</i>	x		x			
<i>Filipendula ulmaria</i>	x	x	x			
<i>Fraxinus excelsior</i>				x		
<i>Fumaria officinalis</i>			x			
<i>Galeopsis tetrahit</i>	x	x	x			
<i>Galium aparine</i>	x	x	x	x		x
<i>Galium mollugo</i>	x	x	x		x	x
<i>Galium palustre</i>			x			
<i>Galium verum</i>					x	x
<i>Geranium molle</i>					x	x

<i>Geranium robertianum</i>		x	x	x	x
<i>Geranium sylvaticum</i>		x	x		
<i>Geum urbanum</i>	x	x	x		x
<i>Glechoma hederacea</i>		x	x	x	
<i>Glyceria maxima</i>	x	x	x		
<i>Hedera helix</i>			x		x
<i>Heracleum sphondylium</i>		x	x		
<i>Hieracium umbellatum</i>				x	x
<i>Himantoglossum hircinum</i>					x
<i>Hippophae rhamnoides</i>				x	x
<i>Holcus lanatus</i>	x	x	x	x	x
<i>Honckenya peploides</i>					x
<i>Humulus lupulus</i>	x	x	x		
<i>Hypericum perforatum</i>		x	x		x
<i>Hypochaeris radicata</i>				x	
<i>Impatiens noli-tangere</i>		x	x		
<i>Impatiens parviflora</i>		x			
<i>Inula conyzoides</i>					x
<i>Jacobaea vulgaris</i>				x	x
<i>Juncus bufonius</i>			x		
<i>Juncus effusus</i>		x	x		
<i>Koeleria macrantha</i>					x
<i>Lamium album</i>		x	x		
<i>Lamium galeobdolon</i>	x		x		
<i>Lapsana communis</i>		x	x		
<i>Lathyrus pratensis</i>		x			
<i>Leontodon saxatilis</i>					x
<i>Lepidium draba</i>				x	x
<i>Ligustrum vulgare</i>					x
<i>Linaria repens</i>			x		
<i>Lolium multiflorum</i>				x	
<i>Lolium perenne</i>	x	x	x		x
<i>Lonicera periclymenum</i>				x	
<i>Lotus corniculatus</i>		x			x
<i>Luzula campestris</i>					x
<i>Lysimachia nummularia</i>			x		
<i>Lysimachia vulgaris</i>	x	x			
<i>Lythrum salicaria</i>	x		x		
<i>Melilotus albus</i>	x		x		
<i>Milium effusum</i>		x	x		
<i>Myosotis arvensis</i>		x	x		
<i>Myosotis ramosissima</i>				x	x
<i>Myosoton aquaticum</i>	x	x			
<i>Oenothera glazioviana</i>				x	x
<i>Ononis repens</i>				x	x

<i>Persicaria amphibia</i>	x	x				
<i>Persicaria hydropiper</i>	x		x			
<i>Persicaria maculosa</i>		x	x			
<i>Phalaris arundinacea</i>	x	x	x			
<i>Phleum arenarium</i>				x	x	
<i>Phleum pratense</i>	x	x	x	x		x
<i>Phragmites australis</i>	x	x	x			
<i>Plantago lanceolata</i>		x		x	x	x
<i>Plantago major</i>		x	x			
<i>Poa pratensis</i>		x	x		x	x
<i>Populus tremula</i>			x			
<i>Potentilla anserina</i>	x	x				
<i>Potentilla reptans</i>		x		x	x	x
<i>Prunella vulgaris</i>		x				
<i>Prunus serotina</i>					x	
<i>Prunus spinosa</i>					x	x
<i>Quercus robur</i>		x			x	
<i>Ranunculus repens</i>	x		x			
<i>Rhinanthus minor</i>				x	x	x
<i>Ribes rubrum</i>		x	x			
<i>Rorippa palustris</i>			x			
<i>Rosa canina</i>					x	x
<i>Rubus caesius</i>				x	x	x
<i>Rubus fruticosus</i>	x	x	x			
<i>Rubus idaeus</i>	x	x	x			
<i>Rumex acetosa</i>		x	x			
<i>Rumex crispus</i>	x	x	x	x	x	
<i>Rumex obtusifolius</i>	x	x	x			
<i>Salix alba</i>				x	x	x
<i>Salix caprea</i>			x			
<i>Salix repens</i>						x
<i>Sambucus nigra</i>			x			
<i>Scrophularia nodosa</i>						
<i>Sedum acre</i>					x	x
<i>Senecio inaequidens</i>				x	x	x
<i>Senecio vulgaris</i>					x	x
<i>Silene dioica</i>	x	x				
<i>Silene latifolia</i>					x	x
<i>Solanum dulcamara</i>		x			x	x
<i>Solidago virgaurea</i>			x			
<i>Sonchus arvensis</i>					x	x
<i>Sonchus asper</i>					x	x
<i>Sonchus oleraceus</i>			x		x	x
<i>Sparganium emersum</i>			x			
<i>Spergula arvensis</i>			x			

<i>Stachys palustris</i>	x	x	x		
<i>Stachys sylvestris</i>	x	x	x		
<i>Stellaria graminea</i>		x	x		
<i>Stellaria holostea</i>		x	x		
<i>Stellaria media</i>		x	x		x
<i>Stellaria nemorum</i>	x		x		
<i>Symphytum officinale</i>	x	x	x		
<i>Tanacetum vulgare</i>					x
<i>Taraxacum officinale</i>	x	x			
<i>Thalictrum minus</i>			x	x	
<i>Tragopogon pratensis</i>				x	x
<i>Trifolium pratense</i>		x	x		
<i>Trifolium repens</i>			x		
<i>Tripleurospermum maritimum</i>			x		
<i>Tussilago farfara</i>		x	x		
<i>Typha latifolia</i>		x			
<i>Urtica dioica</i>	x	x	x	x	x
<i>Valeriana officinalis</i>		x	x		
<i>Valerianella locusta</i>				x	x
<i>Veronica arvensis</i>				x	x
<i>Veronica chamaedrys</i>			x		
<i>Veronica hederifolia</i>				x	
<i>Vicia cracca</i>	x	x			
<i>Vicia hirsuta</i>	x			x	
<i>Vicia sativa</i>			x		x
<i>Viola arvensis</i>		x			

**Appendix 2. Correlation matrix for all measured functional traits. Pearson R values given for the *I. glandulifera* dataset in the lower left triangle, Pearson R values given for the *R. rugosa* dataset in the upper right triangle.** LDMC = leaf dry matter content, LNC = leaf nitrogen content, LPC = leaf phosphorous content, SLA = specific leaf area, SSD = specific stem density. <sup>i</sup>= square root transformed for *I. glandulifera* dataset, <sup>r</sup>= square root transformed for *R. rugosa* dataset.

	Plant height <sup>r</sup>	Leaf area <sup>ir</sup>	SLA <sup>ir</sup>	LDMC <sup>r</sup>	SSD <sup>i</sup>	LNC	LPC
Plant height <sup>r</sup>	1	0.003	-0.274	0.334	-	-0.075	-0.2
Leaf area <sup>ir</sup>	0.214	1	-0.028	0.074	-	-0.152	0.229
SLA <sup>ir</sup>	-0.126	-0.019	1	-0.64	-	0.513	0.414
LDMC <sup>r</sup>	0.174	0.02	-0.686	1	-	-0.421	-0.487
SSD <sup>i</sup>	0.213	-0.028	-0.573	0.746	1	-	-
LNC	-	-	-	-	-	1	0.372
LPC	-	-	-	-	-	-	1

### Appendix 3. Complete reference list for functional traits obtained from the TRY database.

Dataset	Reference
Functional traits explaining variation in plant life history strategies	Adler PB, R Salguero-Gómez, A Compagnoni, JS Hsu, J Ray-Mukherjee, C Mbeau-Ache, M Franco (2014) Functional traits explain variation in plant life history strategies. PNAS 111 (2) 740-745.
Canopy Traits for Temperate Tree Species Under High N-Deposition	Adriaenssens S. (2012). Dry deposition and canopy exchange for temperate tree species under high nitrogen deposition. PhD thesis, Ghent University, Ghent, Belgium, 209p. Atkin OK, KJ Bloomfield, PB Reich, MG Tjoelker, GP Asner, D Bonal, G Bönisch, et al. (2015) Global variability in leaf respiration among plant functional types in relation to climate and leaf traits. <i>New Phytologist</i> .
Global Respiration Database	Atkin, O. K., M. H. M. Westbeek, M. L. Cambridge, H. Lambers, and T. L. Pons. 1997. Leaf respiration in light and darkness - A comparison of slow- and fast-growing Poa species. <i>Plant Physiology</i> 113:961-965.
Plant Physiology Database	Atkin, O. K., M. Schortemeyer, N. McFarlane, and J. R. Evans. 1999. The response of fast- and slow-growing Acacia species to elevated atmospheric CO <sub>2</sub> : an analysis of the underlying components of relative growth rate. <i>Oecologia</i> 120:544-554.
Plant Physiology Database	Blonder, B., Buzzard, B., Sloat, L., Simova, I., Lipson, R., Boyle, B., Enquist, B. (2012) The shrinkage effect biases estimates of paleoclimate. <i>American Journal of Botany</i> . 99.11 1756-1763.
Leaf Structure, Venation and Economic Spectrum	Blonder, B., Vasseur, F., Violle, C., Shipley, B., Enquist, B., Vile, D. <i>Arabidopsis thaliana</i> rejects theories for the origin of the leaf economics spectrum. (in review, <i>New Phytologist</i> )
Leaf Structure, Venation and Economic Spectrum	Blonder, B., Violle, C. and Enquist, B. J. (2013) Assessing the causes and scales of the leaf economics spectrum using venation networks in <i>Populus tremuloides</i> . <i>Journal of Ecology</i> 101: 981–989.
Leaf Structure, Venation and Economic Spectrum	Blonder, B., Violle, C., Patrick, L., Enquist, B. Leaf venation networks and the origin of the leaf economics spectrum. <i>Ecology Letters</i> , 2011.
Italian Alps Plant Traits Database	Bragazza L (2009) Conservation priority of Italian alpine habitats: a floristic approach based on potential distribution of vascular plant species. <i>Biodiversity and Conservation</i> 18: 2823–2835.
Xylem Functional Traits (XFT) Database	Brendan Choat, Steven Jansen et al. (2012) Global convergence in the vulnerability of forests to drought. <i>Nature</i> 491, 752–755.
Xylem Functional Traits (XFT) Database	Brendan Choat, Steven Jansen, Tim J. Brodribb, Herve Cochard, Sylvain Delzon, Radika Bhaskar, Sandra J. Bucci, et al. (2012) Global convergence in the vulnerability of forests to drought. <i>Nature</i> 491:752-755.
Plant Traits from Circeo National Park, Italy	Burrascano S, Copiz R, Del Vico E, Fagiani S, Giarrizzo E, Mei M, Mortelliti A, Sabatini FM, Blasi C (2015) Wild boar rooting intensity determines shifts in understorey composition and functional traits. <i>COMMUNITY ECOLOGY</i> 16(2) 244-253.

Plant Physiology Database

Leaf Traits in Central Apennines Beech Forests

Photosynthetic Capacity Dataset

Sheffield & Spain Woody Database

Sheffield & Spain Woody Database

Floridian Leaf Traits Database

Global Wood Density Database

Mediterranean psammophytes

Plant Traits from Romania

Sheffield Database

Sheffield & Spain Woody Database

Sheffield Database

Abisko & Sheffield Database

- Campbell, C., L. Atkinson, J. Zaragoza-Castells, M. Lundmark, O. Atkin, and V. Hurry. 2007. Acclimation of photosynthesis and respiration is asynchronous in response to changes in temperature regardless of plant functional group. *New Phytologist* 176:375-389.
- Campetella, G.; Botta-Dukát, Z.; Wellstein, C.; Canullo, R.; Gatto, S.; Chelli, S.; Mucina, L.; Bartha, S. (2011): Patterns of plant trait-environment relationships along a forest succession chronosequence. *Agriculture, Ecosystems & Environment*, 145(1), 38-48.
- Carswell, F. E., Meir, P., Wandelli, E. V., Bonates, L. C. M., Kruijt, B., Barbosa, E. M., Nobre, A. D. & Jarvis, P. G. 2000 Photosynthetic capacity in a central Amazonian rain forest. *Tree physiology*. 20, 3, p. 179-186 8 p.
- Castro-Diez, P., J. P. Puyravaud, and J. H. C. Cornelissen. 2000. Leaf structure and anatomy as related to leaf mass per area variation in seedlings of a wide range of woody plant species and types. *Oecologia* 124:476-486.
- Castro-Diez, P., J. P. Puyravaud, J. H. C. Cornelissen, and P. Villar-Salvador. 1998. Stem anatomy and relative growth rate in seedlings of a wide range of woody plant species and types. *Oecologia* 116:57-66.
- Cavender-Bares, J., A. Keen, and B. Miles. 2006. Phylogenetic structure of floridian plant communities depends on taxonomic and spatial scale. *Ecology* 87:S109-S122.
- Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a world wide wood economics spectrum. *Ecology Letters* 12:351-366.
- Ciccarelli D. (2015) - Mediterranean coastal dune vegetation: Are disturbance and stress the key selective forces that drive the psammophilous succession? *Estuarine, Coastal and Shelf Science* 165(5):247–253.
- Ciocarlan V. (2009). The illustrated Flora of Romania. *Pteridophyta et Spermatoptya*. Editura Ceres, 1141 p (in Romanian).
- Cornelissen, J. H. C. 1996. An experimental comparison of leaf decomposition rates in a wide range of temperate plant species and types. *Journal of Ecology* 84:573-582.
- Cornelissen, J. H. C., B. Cerabolini, P. Castro-Diez, P. Villar-Salvador, G. Montserrat-Martí, J. P. Puyravaud, M. Maestro, M. J. A. Werger, and R. Aerts. 2003. Functional traits of woody plants: correspondence of species rankings between field adults and laboratory-grown seedlings? *Journal of Vegetation Science* 14:311-322.
- Cornelissen, J. H. C., B. Cerabolini, P. Castro-Diez, P. Villar-Salvador, G. Montserrat-Martí, J. P. Puyravaud, M. Maestro, M. J. A. Werger, and R. Aerts. 2003. Functional traits of woody plants: correspondence of species rankings between field adults and laboratory-grown seedlings? *Journal of Vegetation Science* 14:311-322.
- Cornelissen, J. H. C., H. M. Quested, D. Gwynn-Jones, R. S. P. Van Logtestijn, M. A. H. De Beus, A. Kondratchuk, T. V. Callaghan, and R. Aerts. 2004. Leaf digestibility and litter decomposability are related in a wide range of subarctic plant species and types. *Functional Ecology* 18:779-786.

Abisko & Sheffield Database

Sheffield Database

Abisko & Sheffield Database

Sheffield Database

Sheffield Database

Sheffield & Spain Woody Database

Plant Traits for Grassland Species (Konza Prairie, Kansas, USA)

Plant Traits for Grassland Species (Konza Prairie, Kansas, USA)

Plant Traits for Grassland Species (Konza Prairie, Kansas, USA)

Global 15N Database

Roots Of the World (ROW) Database

Jasper Ridge leaf chemistry data

Italian Alps Plant Traits Database

Leaf N-Retention Database

Cornelissen, J. H. C., M. J. A. Werger, P. CastroDiez, J. W. A. vanRheenen, and A. P. Rowland. 1997. Foliar nutrients in relation to growth, allocation and leaf traits in seedlings of a wide range of woody plant species and types. *Oecologia* 111:460-469.

Cornelissen, J. H. C., N. Perez-Harguindeguy, S. Diaz, J. P. Grime, B. Marzano, M. Cabido, F. Vendramini, and B. Cerabolini. 1999. Leaf structure and defence control litter decomposition rate across species and life forms in regional floras on two continents. *New Phytologist* 143:191-200.

Cornelissen, J. H. C., P. C. Diez, and R. Hunt. 1996. Seedling growth, allocation and leaf attributes in a wide range of woody plant species and types. *Journal of Ecology* 84:755-765.

Cornelissen, J. H. C., P. C. Diez, and R. Hunt. 1996. Seedling growth, allocation and leaf attributes in a wide range of woody plant species and types. *Journal of Ecology* 84:755-765.

Cornelissen, J. H. C., R. Aerts, B. Cerabolini, M. J. A. Werger, and M. G. A. van der Heijden. 2001. Carbon cycling traits of plant species are linked with mycorrhizal strategy. *Oecologia* 129:611-619.

Cornelissen, J.H.C. 1999. A triangular relationship between leaf size and seed size among woody species: allometry, ontogeny, ecology and taxonomy. *Oecologia* 118: 248-255.

Craine JM, Nippert JB, Towne EG, Tucker S, Kembel SW, Skibbe A, McLaughlan KK (2011) Functional consequences of climate-change induced plant species loss in a tallgrass prairie. *Oecologia* 165: 1109-1117

Craine JM, Ocheltree TW, Nippert JB, Towne EG, Skibbe AM, Kembel SW, Fargione JE (2012) Global diversity of drought tolerance and grassland climate-change resilience. *Nature Climate Change*

Craine JM, Towne EG, Ocheltree TW, Nippert JB (2012) Community traitscape of foliar nitrogen isotopes reveals N availability patterns in a tallgrass prairie. *Plant Soil* 356: 395-403

Craine, J. M., A. J. Elmore, M. P. M. Aidar, M. Bustamante, T. E. Dawson, E. A. Hobbie, A. Kahmen, M. C. Mack, K. K. McLaughlan, A. et al. 2009. Global patterns of foliar nitrogen isotopes and their relationships with climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New Phytologist* 183:980-992.

Craine, J. M., W. G. Lee, W. J. Bond, R. J. Williams, and L. C. Johnson. 2005. Environmental constraints on a global relationship among leaf and root traits of grasses. *Ecology* 86:12-19.

Dahlin KM, Asner GP & CB Field (2013) Environmental and community controls on plant canopy chemistry in a Mediterranean-type ecosystem. *Proceedings of the National Academy of Sciences USA*. 110(17): 6895-6900

Dainese M, Bragazza L (2012) Plant traits across different habitats of the Italian Alps: a comparative analysis between native and alien species. *Alpine Botany* 122: 11-21.

De Vries F., Bardgett R.D. (2016) Plant community controls on short-term ecosystem nitrogen retention. *New Phytologist*.

Sheffield Database

SLA responses to environmental gradients through space and time

The DIRECT Plant Trait Database

Ecological Flora of the British Isles

Fonseca/Wright New South Wales Database

Traits from Subarctic Plant Species Database

The DIRECT Plant Trait Database

BASECO: a floristic and ecological database of Mediterranean French flora

Species and trait shifts in Apennine grasslands

Leaf traits data (SLA) for 56 woody species at the Smithsonian Conservation Biology Institute-Forest

French Alps Trait Data

PLANTSdata USDA

PLANTATT - Attributes of British and Irish Plants

Leaf Physiology Database

Díaz, S., J. G. Hodgson, K. Thompson, M. Cabido, J. H. C. Cornelissen, A. Jalili, G. Montserrat-Martí, J., et al. The plant traits that drive ecosystems: Evidence from three continents. *Journal of Vegetation Science* 15:295-304.

Dwyer, J. M., R. J. Hobbs, and M. M. Mayfield. 2014. Specific leaf area responses to environmental gradients through space and time. *Ecology* 95:399-410

Everwand G, Fry, EL, Eggers T, Manning P (2014) Seasonal variation in the relationship between plant traits and grassland carbon and water fluxes. *Ecosystems* 17, 1095-1108

Fitter, A. H. and H. J. Peat 1994. The Ecological Flora Database. *Journal of Ecology* 82:415-425.

Fonseca, C. R., J. M. Overton, B. Collins, and M. Westoby. 2000. Shifts in trait-combinations along rainfall and phosphorus gradients. *Journal of Ecology* 88:964-977.

Freschet, G. T., J. H. C. Cornelissen, R. S. P. van Logtestijn, and R. Aerts. 2010. Evidence of the 'plant economics spectrum' in a subarctic flora. *Journal of Ecology* 98:362-373.

Fry, E.L., Power, S.A. Manning, P. (2014) Trait based classification and manipulation of functional groups in biodiversity-ecosystem function experiments. *Journal of Vegetation Science*, 25, 248-261.

Gachet S, Errol Véla, Thierry Tatoni, 2005, BASECO: a floristic and ecological database of Mediterranean French flora. *Biodiversity and Conservation* 14(4):1023-1034

Giarrizzo E., Burrascano S., Chiti T., de Bello F., Leps J., Zavattero L., Blasi C. (2017) Re-visiting historical semi-natural grasslands in the Apennines to assess patterns of changes in plant species composition and functional traits. *Applied Vegetation Science* 20(2):247-258.

Gonzalez-Akre, E., McShea, W., Bourg, N., Anderson-Teixeira, K. 2015. Leaf traits data (SLA) for 56 woody species at the Smithsonian Conservation Biology Institute-ForestGEO Forest Dynamic Plot. Front Royal, Virginia. USA. [Data set]. Version 1.0.([www.try-db.org](http://www.try-db.org))

Gos P., Loucugaray G., Colace MP., Arnoldi C., Gaucherand S., Dumazel D., Girard L., Delorme S., Lavorel S. (2016) *Oecologia* 180:1001.

Green, W. 2009. USDA PLANTS Compilation, version 1, 09-02-02.

(<http://bricol.net/downloads/data/PLANTSdatabase/>) NRCS: The PLANTS Database (<http://plants.usda.gov>, 1 Feb 2009). National Plant Data Center: Baton Rouge, LA 70874-74490 USA.

HILL, M.O., PRESTON, C.D. & ROY, D.B. (2004) PLANTATT - attributes of British and Irish Plants: status, size, life history, geography and habitats. Huntingdon: Centre for Ecology and Hydrology.

Kattge, J., W. Knorr, T. Raddatz, and C. Wirth. 2009. Quantifying photosynthetic capacity and its relationship to leaf nitrogen content for global-scale terrestrial biosphere models. *Global Change Biology* 15:976-991.

Leaf and Whole Plant Traits Database

The Americas N&P database

Leaf Traits Mount Hutt, New Zealand

The LEDA Traitbase

Plant traits of grassland species

Plant Traits for Pinus and Juniperus Forests in Arizona

Plant Traits for Pinus and Juniperus Forests in Arizona

French Alps Trait Data

Traits of the Hungarian flora

French Massif Central Grassland Trait Database

Plant Physiology Database

Structural and biochemical leaf traits of boreal tree species in Finland

Photosynthesis Traits Worldwide

Kazakou, E., D. Vile, B. Shipley, C. Gallet, and E. Garnier. 2006. Co-variations in litter decomposition, leaf traits and plant growth in species from a Mediterranean old-field succession. *Functional Ecology* 20:21-30.

Kerkhoff, A. J., W. F. Fagan, J. J. Elser, and B. J. Enquist. 2006. Phylogenetic and growth form variation in the scaling of nitrogen and phosphorus in the seed plants. *American Naturalist* 168:E103-E122.

Kichenin et al. 2013. Contrasting effects of plant inter- and intraspecific variation on community-level trait measures along an environmental gradient. *Functional Ecology*, in press.

Kleyer, M., R. M. Bekker, I. C. Knevel, J. P. Bakker, K. Thompson, M. Sonnenschein, P. Poschlod, J., et al. 2008. The LEDA Traitbase: a database of life-history traits of the Northwest European flora. *Journal of Ecology* 96:1266-1274.

La Pierre, KJ and Smith, MD. (2015) Functional trait expression of grassland species shift with short- and long-term nutrient additions. *Plant Ecology* 216: 307.

Laughlin, D. C., J. J. Leppert, M. M. Moore, and C. H. Sieg. 2010. A multi-trait test of the leaf-height-seed plant strategy scheme with 133 species from a pine forest flora. *Functional Ecology* 24:493-501.

Laughlin, D.C., P.Z. Fulé, D.W. Huffman, J. Crouse, and E. Laliberte. 2011. Climatic constraints on trait-based forest assembly. *Journal of Ecology* 99:1489-1499.

Lavorel, S., Grigulis, K., Lamarque, P., Colace, M.-P., Garden, D., Girel, J., Pellet, G. and Douzet, R. (2011), Using plant functional traits to understand the landscape distribution of multiple ecosystem services. *Journal of Ecology*, 99: 135–147.

Lhotsky B., Anikó Csecserits, Bence Kovács, Zoltán Botta-Dukát: New plant trait records of the Hungarian flora.

Louault, F., V. D. Pillar, J. Aufreire, E. Garnier, and J. F. Soussana. 2005. Plant traits and functional types in response to reduced disturbance in a semi-natural grassland. *Journal of Vegetation Science* 16:151-160.

Loveys, B. R., L. J. Atkinson, D. J. Sherlock, R. L. Roberts, A. H. Fitter, and O. K. Atkin. 2003. Thermal acclimation of leaf and root respiration: an investigation comparing inherently fast- and slow-growing plant species. *Global Change Biology* 9:895-910.

Lukeš, P., Stenberg, P., Rautiainen, M., Möttus, M., Vanhatalo, K.M. Optical properties of leaves and needles for boreal tree species in Europe (2013) *Remote Sensing Letters*, 4 (7), pp. 667-676

Maire V, Ian J. Wright, I. Colin Prentice, Niels H. Batjes, Radika Bhaskar, Peter M. van Bodegom, Will K. Cornwell, David Ellsworth, Ülo Niinemets, Alejandro Ordoñez, Peter B. Reich, Louis S. Santiago (2015). Global soil and climate effects on leaf photosynthetic traits and rates. *Global Ecology and Biogeography* 24(6): 706-717. Maire V, Wright IJ, Prentice IC, Batjes NH, Bhaskar R, van Bodegom PM, Cornwell WK, Ellsworth D, Niinemets Ü, Ordoñez A, Reich PB, Santiago LS (2015) Data from:

Fonseca/Wright New South Wales Database

Leaf and Whole Plant Traits Database

ECOCRAFT

ECOCRAFT

ECOCRAFT

Photosynthetic Capacity Dataset

Photosynthetic Capacity Dataset

Whole Plant Hydraulic Conductance

Chinese Traits

Leaf and Whole Plant Traits Database

Leaf and Whole Plant Traits Database

Altitudinal Vicariants Spain

Traits of halophytic species

Global effects of soil and climate on leaf photosynthetic traits and rates. Dryad Digital Repository.  
<http://dx.doi.org/10.5061/dryad.j42m7>

McDonald, P. G., C. R. Fonseca, J. M. Overton, and M. Westoby. 2003. Leaf-size divergence along rainfall and soil-nutrient gradients: is the method of size reduction common among clades? *Functional Ecology* 17:50-57.

McKenna, M. F. and B. Shipley. 1999. Interacting determinants of interspecific relative growth: Empirical patterns and a theoretical explanation. *Ecoscience* 6:286-296.

Medlyn, B. E. and P. G. Jarvis. 1999. Design and use of a database of model parameters from elevated [CO<sub>2</sub>] experiments. *Ecological Modelling* 124:69-83.

Medlyn, B. E., C. V. M. Barton, M. S. J. Broadmeadow, R. Ceulemans, P. DeAngelis, M. Forstreuter, M. Freeman, S. B. Jackson, S. Kellomaeki, E. Laitat, A. Rey, P. Roberntz, B. D. Sigurdsson, J.

Strassemeyer, K. Wang, P. S. Curtis, and P. G. Jarvis. 2001. Stomatal Conductance of forest species after long-term exposure to elevated CO<sub>2</sub> concentration: a synthesis. *New Phytologist* 149:247--264.

Medlyn, B. E., F.-W. Badeck, D. G. G. De Pury, C. V. M. Barton, M. Broadmeadow, R. Ceulemans, P. De Angelis, M. Forstreuter, M. E., et al. 1999. Effects of elevated CO<sub>2</sub> on photosynthesis in European forest species: a meta-analysis of model parameters. *Plant, Cell and Environment* 22:1475-1495.

Meir, P. & Levy, P. E. 2007 Photosynthetic parameters from two contrasting woody vegetation types in West Africa. *Plant Ecology*. 192, 2, p. 277-287 11 p.

Meir, P., Kruijt, B., Broadmeadow, M., Kull, O., Carswell, F. & Nobre, A. 2002 Acclimation of photosynthetic capacity to irradiance in tree canopies in relation to leaf nitrogen concentration and leaf mass per unit area. *Plant, Cell and Environment*. 25, 3, p. 343-357 15 p.

MENCUCCINI M., 2003. The ecological significance of long distance water transport: short-term regulation and long-term acclimation across plant growth forms. *Plant, Cell and Environment*, 26:163-182.

Meng, T.-T., Wang, H., Harrison, S. P., Prentice, I. C., Ni, J., and Wang, G.: Responses of leaf traits to climatic gradients: adaptive variation versus compositional shifts, *Biogeosciences*, 12, 5339-5352, 2015.

Meziane, D. and B. Shipley. 1999. Interacting components of interspecific relative growth rate: constancy and change under differing conditions of light and nutrient supply. *Functional Ecology* 13:611-622.

Meziane, D. and B. Shipley. 1999. Interacting determinants of specific leaf area in 22 herbaceous species: effects of irradiance and nutrient availability. *Plant Cell and Environment* 22:447-459.

Milla & Reich 2011 *Annals of Botany* 107: 455–465, 2011.

Minden V, M Kleyer (2015): Ecosystem multifunctionality of coastal marshes is determined by key plant traits, *Journal of Vegetation Science* 26: 651-662

- Element contents of plant organs of halohytic species, NW-Germany  
Minden V, Michael Kleyer (2014): Internal and external regulation of plant organ stoichiometry, *Plant Biology*, 16: 897-907
- Traits of Halohytic Species in North-West-Germany  
Minden Vanessa, Michael Kleyer (2011): Testing the effect–response framework: key response and effect traits determining above-ground biomass of salt marshes. *Journal of Vegetation Science* 22: 387-401
- The Netherlands Plant Traits Database  
Ordonez, J. C., P. M. van Bodegom, J. P. M. Witte, R. P. Bartholomeus, H. F. van Dobben, and R. Aerts. 2010. Leaf habit and woodiness regulate different leaf economy traits at a given nutrient supply. *Ecology* 91:3218-3228.
- The Netherlands Plant Traits Database  
Ordonez, J. C., P. M. van Bodegom, J. P. M. Witte, R. P. Bartholomeus, J. R. van Hal, and R. Aerts. 2010. Plant Strategies in Relation to Resource Supply in Mesic to Wet Environments: Does Theory Mirror Nature? *American Naturalist* 175:225-239.
- Impatiens Glandulifera Dataset  
Pahl, A.T., Kollmann, J., Mayer, A. & Haider, S. (2013): No evidence for local adaptation in an invasive alien plant: field and greenhouse experiments tracing a colonization sequence. *Annals of Botany* 112 (9): 1921-1930.
- BROT Plant Trait Database  
Paula, S. and J. G. Pausas. 2008. Burning seeds: germinative response to heat treatments in relation to resprouting ability. *Journal of Ecology* 96:543-552.
- BROT Plant Trait Database  
Paula, S., M. Arianoutsou, D. Kazanis, Ç. Tavsanoglu, F. Lloret, C. Buhk, F. Ojeda, B. Luna, J. M. Moreno, A. Rodrigo, J. M. Espelta, S. Palacio, B. Fernández-Santos, P. M. Fernandes, and J. G. Pausas. 2009. Fire-related traits for plant species of the Mediterranean Basin. *Ecology* 90:1420.
- Plant Traits of Acidic Grasslands in Central Spain  
Peco B., de Pablos I., Traba J. , & Levassor C. (2005) The effect of grazing abandonment on species composition and functional traits: the case of dehesa *Basic and Applied Ecology*, 6(2): 175-183
- Chinese Traits  
Prentice, I.C., Meng, T., Wang, H., Harrison, S.P., Ni, J., Wang, G., 2011. Evidence for a universal scaling relationship of leaf CO<sub>2</sub> drawdown along a moisture gradient. *New Phytologist* 190: 169–180
- Leaf and Whole Plant Traits Database  
Pyankov, V. I., A. V. Kondratchuk, and B. Shipley. 1999. Leaf structure and specific leaf mass: the alpine desert plants of the Eastern Pamirs, Tadzhikistan. *New Phytologist* 143:131-142.
- Abisko & Sheffield Database  
Quested, H. M., J. H. C. Cornelissen, M. C. Press, T. V. Callaghan, R. Aerts, F. Trosien, P. Riemann, D. Gwynn-Jones, A. Kondratchuk, and S. E. Jonasson. 2003. Decomposition of sub-arctic plants with differing nitrogen economies: A functional role for hemiparasites. *Ecology* 84:3209-3221.
- Plant Traits from Romania  
Sanda V., Bita-Nicolae C.D. & Barabas N. (2003). The flora of spontane and cultivated cormophytes from Romania. Editura “Ion Borcea”, Bacau, 316 p (in Romanian)
- California Coastal Grassland Database  
Sandel, B., J. D. Corbin, and M. Krupa. 2011. Using plant functional traits to guide restoration: a case study in California coastal grassland. *Ecosphere* 2(2):art23.
- BIOTREE Trait Data  
Scherer-Lorenzen, M., Schulze, E.-D., Don, A., Schumacher, J. & Weller, E. (2007) Exploring the functional significance of forest diversity: A new long-term experiment with temperate tree species (BIOTREE). *Perspectives in Plant Ecology, Evolution and Systematics*, 9, 53-70.

Traits of 59 grassland species  
The Xylem/Phloem Database  
The Xylem/Phloem Database  
Herbs Water Relations on Soil Moisture Gradients  
Leaf and Whole Plant Traits Database  
Eastern US Old Field Plant Traits Database  
Old fields of Eastern US (Siefert Data)  
Root Traits of Grassland Species  
Niwot Alpine Plant Traits  
Grassland Plant Trait Database  
Grassland Plant Trait Database

Schroeder-Georgi, T., Wirth, C., Nadrowski, K., Meyer, S. T., Mommer, L. and Weigelt, A. (2016), From pots to plots: hierarchical trait-based prediction of plant performance in a mesic grassland. *J Ecol*, 104: 206–218.

Schweingruber, F.H., Landolt, W.: The Xylem Database. Swiss Federal Research Institute WSL Updated (2005).

Schweingruber, F.H., Poschlod, P. 2005: Growth rings in herbs and shrubs: Life span, age determination and stem anatomy. *Forest, Snow and Landscape Res.* 79, 195-415.

Sheremetev S.N. (2005) Herbs on the soil moisture gradient (water relations and the structural-functional organization). KMK, Moscow, 271 pp. (In Russian).

Shipley B., 2002. Trade-offs between net assimilation rate and specific leaf area in determining relative growth rate: relationship with daily irradiance, *Functional Ecology*(16) 682-689.

Shipley, B. 1989. The Use of above-Ground Maximum Relative Growth-Rate as an Accurate Predictor of Whole-Plant Maximum Relative Growth-Rate. *Functional Ecology* 3:771-775.

Shipley, B. 1995. Structured Interspecific Determinants of Specific Leaf-Area in 34 Species of Herbaceous Angiosperms. *Functional Ecology* 9:312-319.

Shipley, B. and M. J. Lechowicz. 2000. The functional co-ordination of leaf morphology, nitrogen concentration, and gas exchange in 40 wetland species. *Ecoscience* 7:183-194.

Shipley, B. and M. Parent. 1991. Germination Responses of 64 Wetland Species in Relation to Seed Size, Minimum Time to Reproduction and Seedling Relative Growth-Rate. *Functional Ecology* 5:111-118.

Shipley, B. and T. T. Vu. 2002. Dry matter content as a measure of dry matter concentration in plants and their parts. *New Phytologist* 153:359-364.

Siefert, A., Fridley, J.D., and Ritchie, M.E. 2014. Community functional responses to soil and climate at multiple spatial scales: when does intraspecific variation matter? *PLOS ONE* 9: e111189

Siefert, A., Fridley, J.D., and Ritchie, M.E. 2014. Community functional responses to soil and climate at multiple spatial scales: when does intraspecific variation matter? *PLOS ONE* 9: e111189

Smith, S. W., Woodin, S. J., Pakeman, R. J., Johnson, D. and van der Wal, R. (2014), Root traits predict decomposition across a landscape-scale grazing experiment. *New Phytologist*.

Spasojevic, M. J. and K. N. Suding. 2012. Inferring community assembly mechanisms from functional diversity patterns: the importance of multiple assembly processes. *Journal of Ecology* 100:652-661.

Takkis, K. 2014. Changes in plant species richness and population performance in response to habitat loss and fragmentation. *DISSERTATIONES BIOLOGICAE UNIVERSITATIS TARTUENSIS* 255, 2014-04-07. Available from: <http://hdl.handle.net/10062/39546>

Takkis, K., Saar, L., Pärtel, M., Helm, A. Effect of environment and landscape on the traits of six plant species in fragmented grasslands. (in preparation)

- Leaf Traits and Seed Mass of Cover Crops
- Plant Traits for Grassland Species (Konza Prairie, Kansas, USA)
- Nutrient Resorption Efficiency Database
- Nutrient Resorption Efficiency Database
- Leaf and Whole Plant Traits Database
- A Global Data Set of Leaf Photosynthetic Rates, Leaf N and P, and Specific Leaf Area
- A Global Data Set of Leaf Photosynthetic Rates, Leaf N and P, and Specific Leaf Area
- Global Leaf Element Composition Database
- Leaf Ash Content in China's Terrestrial Plants
- The Functional Ecology of Trees (FET) Database - Jena
- Leaf Economic Traits Across Varying Environmental Conditions
- GLOPNET - Global Plant Trait Network Database
- Tribouillois H, Fort F, Cruz P, Charles R, Flores O, Garnier E, et al. (2015) A Functional Characterisation of a Wide Range of Cover Crop Species: Growth and Nitrogen Acquisition Rates, Leaf Traits and Ecological Strategies. *PLoS ONE* 10(3): e0122156.
- Tucker SS, Craine JM, Nippert JB (2011) Physiological drought tolerance and the structuring of tallgrass assemblages. *Ecosphere* 2(4): 48.
- Vergutz, L., Manzoni, S., Porporato, A., Novais, R. F., and Jackson, R. B. 2012. Global resorption efficiencies and concentrations of carbon and nutrients in leaves of terrestrial plants, *Ecological Monographs* 82:2, 205-220.
- Vergutz, L., S. Manzoni, A. Porporato, R.F. Novais, and R.B. Jackson. 2012. A Global Database of Carbon and Nutrient Concentrations of Green and Senesced Leaves. Data set. Available on-line [<http://daac.ornl.gov>] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. <http://dx.doi.org/10.3334/ORNLDAAAC/1106>
- Vile, D. 2005. Significations fonctionnelle et ecologique des traits des especes vegetales: exemple dans une succession post-cultural mediterraneenne et generalisations, PHD Thesis.
- Walker AP, AP Beckerman, L Gu , J Kattge, LA Cernusak, TF Domingues, JC Scales, G Wohlfahrt, SD Wullschleger and FI Woodward (2014) The relationship of leaf photosynthetic traits –  $V_{cmax}$  and  $J_{max}$  – to leaf nitrogen, leaf phosphorus, and specific leaf area: a meta-analysis and modeling study. *Ecology and Evolution*.
- Walker, A.P. 2014. A Global Data Set of Leaf Photosynthetic Rates, Leaf N and P, and Specific Leaf Area. Data set. Available on-line [<http://daac.ornl.gov>] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA.
- Watanabe, T., M. R. Broadley, S. Jansen, P. J. White, J. Takada, K. Satake, T. Takamatsu, S. J. Tuah, and M. Osaki. 2007. Evolutionary control of leaf element composition in plants. *New Phytologist* 174:516-523.
- Wenxuan Han, Yahan Chen, Fang-Jie Zhao, Luying Tang, Rongfeng Jiang and Fusuo Zhang, 2012, Floral, climatic and soil pH controls on leaf ash content in China's terrestrial plants. *Global Ecology and Biogeography*.
- Wirth, C. and J. W. Lichstein. 2009. The Imprint of Species Turnover on Old-Growth Forest Carbon Balances - Insights From a Trait-Based Model of Forest Dynamics. Pages 81-113 in C. Wirth, G. Gleixner, and M. Heimann, editors. *Old-Growth Forests: Function, Fate and Value*. Springer, New York, Berlin, Heidelberg.
- Wright JP, Sutton-Grier A (2012) Does the leaf economic spectrum hold within local species pools across varying environmental conditions? *Functional Ecology* 2012.
- Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-Bares, T. Chapin, J. H. C. Cornelissen, M. Diemer, J. Flexas, E. Garnier, P. K. Groom, J. Gulias, K. Hikosaka, B.

GLOPNET - Global Plant Trait Network Database

Leaf Nitrogen and Phosphorus for China's Terrestrial Plants

Quercus Leaf C&N Database

Global Wood Density Database

Baccara - Plant Traits of European Forests  
Cold Tolerance, Seed Size and Height of North American Forest Tree Species

French Weeds Trait Database  
Growth and Herbivory of Juvenile Trees

Herbaceous Plants Traits From Southern Germany

Meadow Plant Traits: Biomass Allocation, Rooting depth  
New South Wales Plant Traits Database

Overton/Wright New Zealand Database

Photosynthesis and Leaf Characteristics Database

Plant Coastal Dune Traits (France, Aquitaine)

B. Lamont, T. Lee, W. Lee, C. Lusk, J. J. Midgley, M. L. Navas, U. Niinemets, J. Oleksyn, N. Osada, H. Poorter, P. Poot, L. Prior, V. I. Pyankov, C. Roumet, S. C. Thomas, M. G. Tjoelker, E. J. Veneklaas, and R. Villar. 2004. The worldwide leaf economics spectrum. *Nature* 428:821-827.

Wright, I. J., P. B. Reich, O. K. Atkin, C. H. Lusk, M. G. Tjoelker, and M. Westoby. 2006. Irradiance, temperature and rainfall influence leaf dark respiration in woody plants: evidence from comparisons across 20 sites. *New Phytologist* 169:309-319.

Yahan Chen , Wenxuan Han , Luying Tang , Zhiyao Tang and Jingyun Fang 2011 Leaf nitrogen and phosphorus concentrations of woody plants differ in responses to climate, soil and plant growth form. *Ecography* 34.

Yguel B., Bailey R., Tosh N.D., Vialatte A., Vasseur C., Vitrac X., Jean F. & Prinzing A. (2011). Phytophagy on phylogenetically isolated trees: why hosts should escape their relatives. *Ecol. Lett.*, 14, 1117-1124.

Zanne, A. E., G. Lopez-Gonzalez, D. A. Coomes, J. Ilic, S. Jansen, S. L. Lewis, R. B. Miller, N. G. Swenson, M. C. Wiemann, and J. Chave. 2009 Global wood density database. Dryad:Identifier: <http://hdl.handle.net/10255/dryad.10235>.

unpub.

unpub.

unpub.

unpub.

unpub.

unpub.

unpub.

unpub.


unpub.

Rocky Mountain Biological Laboratory WSR/gradient plant traits	unpub.
Trait Data from Niwot Ridge LTER (2016)	unpub.
Traits for Common Grasses and Herbs in Spain	unpub.
Traits of <i>Hypochaeris radicata</i> under shade and drought conditions	unpub.
Tundra Plant Traits Database	unpub.

---


**Appendix 4. Average community weighted mean (CWM) trait values for the control plots (NAT<sub>C</sub>) and the invaded plots (NAT<sub>I</sub>), and average trait values for the invader (INV) for both datasets.** LDMC = leaf dry matter content, LNC = leaf nitrogen content, LPC = leaf phosphorous content, SLA = specific leaf area, SSD = specific stem density.

	Plant height	Leaf area	SLA	LDMC	SSD	LNC	LPC
<b><i>Impatiens glandulifera</i> dataset</b>							
CWM NAT <sub>C</sub>	112.33	4392.26	27.90	259.83	0.224	-	-
CWM NAT <sub>I</sub>	109.57	4104.44	30.92	245.45	0.211	-	-
INV	146.85	5704.55	39.24	142.40	0.077	-	-
<b><i>Rosa rugosa</i> dataset</b>							
CWM NAT <sub>C</sub>	59.37	2331.89	16.64	306.47	-	22.47	1.94
CWM NAT <sub>I</sub>	59.65	2036.27	24.44	249.36	-	24.99	2.19
INV	64.82	4935.54	10.43	368.87	-	21.15	2.35



#### Appendix 5. Visualization of the trait hypervolume in the four trait (PCA) dimensions for

**the *I. glandulifera* dataset.** Large points depict hypervolume centroids, normal points depict data points, small points depict randomized points. Green = control community (NAT<sub>C</sub>), blue= invaded community (NAT<sub>I</sub>), red= invader (INV, *Impatiens glandulifera*).



#### **Appendix 6. Three dimensional gif visualization of the trait hypervolume in the four trait**

**(PCA) dimensions for the *I. glandulifera* dataset.** Normal points depict data points, small points depict randomized points. Green = control community ( $\text{NAT}_C$ ), blue= invaded community ( $\text{NAT}_I$ ), red= invader (INV, *Impatiens glandulifera*).


**Appendix 7. Size ( $SD^4$ ) of the hypervolumes for the *I. glandulifera* dataset containing native plants in the control plots (NAT<sub>C</sub>), native plants in the invaded plots (NAT<sub>I</sub>) and the *I. glandulifera* records (INV) for the five study regions separately, for the average of the five study regions and for the full dataset across all study regions (all data). Average and 95% confidence intervals given based on a standardized number of samples (N=34). Values on the first row compare hypervolumes that were constructed from the measured traits, values on the second row represent hypervolumes analyses extended with database traits.**

Study region	volume INV	volume NAT <sub>C</sub>	Volume NAT <sub>I</sub>
Ghent		49.1 (44.8-53.4)	47.5 (43.5-51.5)
		51.8 (45.8-57.7)	55.4 (48.7-62.1)
Bremen		48.5 (44.9-52.1)	53.8 (50.9-56.8)
		58.3 (52.5-64.2)	62.7 (57.3-68.1)
Lund		45.3 (41.1-49.5)	49.4 (45.8-53.1)
		49.8 (43.7-55.8)	56.3 (50.8-61.7)
Stockholm		43.7 (39.5-47.9)	49.3 (48.3-50.3)
		53.0 (46.2-59.8)	58.7 (51.9-65.4)
Trondheim		39.4 (36.0-42.7)	41.4 (39.2-43.5)
		51.0 (44.8-57.1)	54.0 (49.1-58.9)
<i>average</i>		<b>45.2 (41.3-49.1)</b>	<b>48.3 (53.0-59.2)</b>
		<b>52.8 (46.6-58.9)</b>	<b>57.4 (51.6-63.3)</b>
<b>ALL</b>	<b>14.4 (10.5-18.4)</b>	<b>50.2 (40.8-59.6)</b>	<b>52.4 (44.5-60.4)</b>
		<b>52.4 (46.2-62.4)</b>	<b>56.8 (47.7-65.8)</b>


**Appendix 8. Pairwise hypervolume overlap metrics between the hypervolume of native plants in the control plots (NAT<sub>C</sub>), native plants in the invaded plots (NAT<sub>I</sub>) and the invader plants (INV) of the *I. glandulifera* dataset for the five study regions separately, for the average of the five study regions and for the full dataset across all study regions (all data).** Average and 95% confidence intervals given for the distance between hypervolume centroids, Jaccard similarity between both hypervolumes and the unique volume fraction of both hypervolumes. Each parameter is based on a standardized number of samples (N=34). Values on the first row for hypervolumes on the measured traits, values on the second row for hypervolumes on measured and database traits.

No TRY	Study region	centroid distance (SD <sup>4</sup> )	Jaccard similarity	Unique volume fraction (%)
NAT <sub>C</sub> - NAT <sub>I</sub>	Ghent	0.312 (0.125-0.498)	0.369 (0.330-0.409)	46.9 (41.6-52.2)/45.1 (40.0-50.2)
		0.323 (0.090-0.556)	0.378 (0.292-0.465)	43.3 (34.1-52.4)/46.9 (36.1-57.6)
	Bremen	0.230 (0.049-0.412)	0.397 (0.368-0.426)	40.0 (36.3-43.8)/46.0 (42.1-49.9)
		0.526 (0.189-0.863)	0.330 (0.268-0.392)	48.5 (40.5-56.6)/52.1 (44.7-59.6)
	Lund	0.421 (0.261-0.580)	0.306 (0.266-0.345)	51.0 (45.1-56.9)/55.2 (50.2-60.1)
		0.453 (0.165-0.741)	0.343 (0.278-0.408)	45.6 (36.9-54.3)/51.9 (43.9-59.9)
	Stockholm	0.490 (0.332-0.649)	0.343 (0.308-0.377)	45.7 (41.0-50.4)/51.9 (47.5-56.3)
		0.507 (0.206-0.807)	0.347 (0.278-0.417)	45.7 (36.0-55.4)/51.0 (42.9-59.0)
	Trondheim	0.577 (0.390-0.763)	0.393 (0.350-0.435)	42.2 (37.3-47.0)/45.0 (39.7-50.3)
		0.472 (0.175-0.769)	0.400 (0.297-0.502)	41.3 (30.6-52.0)/44.6 (32.8-56.4)
<i>average</i>		<b>0.406 (0.232-0.580)</b>	<b>0.361 (0.324-0.398)</b>	<b>45.2 (40.3-50.1)/48.6 (43.9-53.4)</b>
		<b>0.456 (0.165-0.747)</b>	<b>0.360 (0.283-0.437)</b>	<b>44.9 (35.6-54.1)/49.3 (40.1-58.5)</b>
	<i>all data</i>	<b>0.490 (0.150-0.829)</b>	<b>0.313 (0.243-0.384)</b>	<b>51.1 (40.8-61.4)/53.2 (43.6-62.9)</b>
NAT <sub>C</sub> - INV	Ghent	2.065 (1.920-2.209)	0.039 (0.018-0.059)	95.3 (92.8-97.8)/82.9 (73.7-92.0)
		2.154 (1.972-2.336)	0.030 (0.006-0.053)	96.4 (93.6-99.2)/86.0 (75.1-96.9)
	Bremen	2.622 (2.543-2.700)	0.001 (<0.001-0.002)	99.8 (99.7-99.9)/99.4 (98.8-99.9)
		2.702 (2.516-2.888)	0.007 (<0.001-0.017)	99.1 (98.0-100)/95.6 (89.9-100)
	Lund	2.560 (2.431-2.688)	0.009 (0.003-0.016)	98.9 (98.0-99.7)/95.8 (92.8-98.8)
		2.103 (1.930-2.275)	0.008 (<0.001-0.015)	99.0 (98.2-99.9)/96.2 (92.5-99.9)
	Stockholm	1.857 (1.750-1.963)	0.012 (0.001-0.023)	98.4 (97.0-99.9)/95.2 (90.9-99.5)
		2.575 (2.353-2.796)	0.016 (<0.001-0.035)	98.0 (95.7-100)/92.7 (83.9-100)
	Trondheim	2.290 (2.158-2.422)	0.009 (<0.001-0.019)	98.8 (97.6-100)/95.6 (91.0-100)
		2.158 (1.975-2.340)	0.011 (<0.001-0.022)	98.7 (97.4-99.9)/93.6 (87.4-99.8)
<i>average</i>		<b>2.279 (2.161-2.397)</b>	<b>0.014 (0.004-0.024)</b>	<b>98.2 (97.0-99.5)/93.8 (89.4-98.1)</b>
		<b>2.338 (2.149-2.527)</b>	<b>0.014 (&lt;0.001-0.028)</b>	<b>98.2 (96.6-99.9)/92.8 (85.7-99.9)</b>
	<i>all data</i>	<b>2.238 (1.917-2.559)</b>	<b>0.010 (&lt;0.001-0.029)</b>	<b>98.7 (96.3-100)/95.4 (87.6-100)</b>

		<b>2.283 (1.955-2.610)</b>	<b>0.011 (&lt;0.001-0.029)</b>	<b>98.6 (96.4-100)/94.7 (86.8-100)</b>
NAT <sub>I</sub> - INV	Ghent	1.971 (1.831-2.111) 2.062 (1.874-2.250)	0.044 (0.023-0.065) 0.032 (0.006-0.058)	94.6 (92.1-97.1)/81.0 (71.7-90.2) 96.1 (93.1-99.2)/84.0 (71.0-97.0)
	Bremen	2.571 (2.479-2.662) 2.630 (2.467-2.793)	0.001 (<0.001-0.002) <0.001 (<0.001-0.001)	99.9 (99.9-100)/99.9 (99.6-100) 99.9 (99.9-100)/99.7 (99.2-100)
	Lund	1.913 (1.817-2.010) 2.269 (2.059-2.480)	0.012 (0.007-0.018) 0.009 (0.003-0.015)	98.5 (97.8-99.2)/94.1 (91.3-96.8) 98.9 (98.1-99.6)/95.0 (91.7-98.4)
	Stockholm	2.469 (2.444-2.493) 2.578 (2.365-2.791)	0.019 (0.016-0.022) 0.014 (<0.001-0.028)	97.6 (97.2-98.0)/91.7 (90.3-93.1) 98.3 (96.6-100)/93.0 (85.9-100)
	Trondheim	2.026 (1.942-2.110) 2.094 (1.932-2.257)	0.035 (0.022-0.048) 0.025 (0.008-0.041)	95.8 (94.2-97.4)/83.7 (77.7-89.7) 97.1 (95.3-99.0)/85.4 (75.6-95.1)
	<i>average</i>	<b>2.190 (2.103-2.277)</b> <b>2.327 (2.139-2.514)</b>	<b>0.022 (0.013-0.031)</b> <b>0.016 (0.003-0.029)</b>	<b>97.3 (96.3-98.3)/90.0 (86.1-94.0)</b> <b>98.1 (96.6-99.6)/91.4 (84.7-98.2)</b>
	<i>all data</i>	<b>2.167 (1.795-2.539)</b> <b>2.252 (1.907-2.596)</b>	<b>0.016 (&lt;0.001-0.041)</b> <b>0.015 (&lt;0.001-0.037)</b>	<b>97.9 (94.9-100)/92.6 (82.0-100)</b> <b>98.1 (95.5-100)/92.2 (81.1-100)</b>



**Appendix 9. Visualization of the trait hypervolume in the four trait (PCA) dimensions for the *R. rugosa* dataset.** Large points depict hypervolume centroids, normal points depict data points, small points depict randomized points. Green = control community (NATc), blue= invaded community (NATi), red= invader (INV, *Rosa rugosa*).



#### Appendix 10. Three dimensional gif visualization of the trait hypervolume in the four trait

**(PCA) dimensions for the *R. rugosa* dataset.** Normal points depict data points, small points depict randomized points. Green = control community (NAT<sub>C</sub>), blue= invaded community (NAT<sub>I</sub>), red= invader (INV, *Rosa rugosa*).

**Appendix 11. Paired t-tests comparing soil conditions between control plots and invaded plots for *Impatiens glandulifera* and *Rosa rugosa*.** Total nitrogen,  $\text{NO}_3^-$  and  $\text{NH}_4^+$  are soil available nitrogen proxies, which were quantified with PRS probes (Plant Root Simulator, Western Ag Innovations, Saskatoon, Canada) during the growing season. Test-statistic (t) and p-value presented for each paired t-test. SOM = soil organic matter.

	<i>I. glandulifera</i>		<i>R. rugosa</i>	
	t	p	t	p
Total N	0.118	0.907	1.539	0.138
$\text{NO}_3^-$	0.196	0.845	1.512	0.145
$\text{NH}_4^+$	0.887	0.379	1.456	0.158
pH	0.545	0.589	1.224	0.233
SOM	1.935	0.059	-0.9203	0.367